

Two-Loop Superstrings in Hyperelliptic Language III: the Four-Particle Amplitude

Zhu-Jun Zheng*

Institute of Mathematics, Henan University
Kaifeng 475001, P. R. China
and

Institute of Theoretical Physics, Chinese Academy of Sciences
P. O. Box 2735, Beijing 100080, P. R. China

Jun-Bao Wu
School of Physics, Peking University
Beijing 100871, P. R. China

Chuan-Jie Zhu[†]
Institute of Theoretical Physics, Chinese Academy of Sciences
P. O. Box 2735, Beijing 100080, P. R. China

February 1, 2008

*Supported in part by Math. Tianyuan Fund with grant Number 10226002 and the Natural Science Foundation of Educational Committee of Henan Province with grant Number 2000110010

[†]Supported in part by fund from the National Natural Science Foundation of China with grant Number 90103004.

Abstract

We compute explicitly the four-particle amplitude in superstring theories by using the hyperelliptic language and the newly obtained chiral measure of D'Hoker and Phong. Although the algebra of the intermediate steps is a little bit involved, we obtain a quite simple expression for the four-particle amplitude. As expected, the integrand is independent of all the insertion points. As an application of the obtained result, we show that the perturbative correction to the R^4 term in type II superstring theories is vanishing point-wise in (even) moduli space at two loops.

1 Introduction

Although we believe that superstring theory is finite in perturbation at any order [1, 2, 3, 4], a rigorous proof is still lacking despite great advances in the covariant formulation of superstring perturbation theory á la Polyakov. The main problem is the presence of supermoduli and modular invariance in higher genus. At two loops these problems were solved explicitly by using the hyperelliptic formalism in a series of papers [5, 6, 7, 8, 9]. The explicit result was also used by Iengo [10] to prove the vanishing of perturbative correction to the R^4 term [11] at two loop, in agreement with the indirect argument of Green and Gutperle [12], Green, Gutperle and Vanhove [13], and Green and Sethi [14] that the R^4 term does not receive perturbative contributions beyond one loop. Recently, Stieberger and Taylor [15] also used the result of [8] to prove the vanishing of the heterotic two-loop F^4 term. For some closely related works we refer the reader to the reviews [16, 17]. In the general case in superstring perturbation theory, there is no satisfactory solution. For a review of these problem we refer the reader to [18, 19].

Recently two-loop superstring was studied by D'Hoker and Phong. In a series of papers [20, 21, 22, 23] (for a recent review see [19]), D'Hoker and Phong found an unambiguous and slice-independent two-loop superstring measure on moduli space for even spin structure from first principles.

Although their result is quite explicit, it is still a difficult problem to use it in actual computation. In [23], D'Hoker and Phong used their result to compute explicitly the chiral measure by choosing the split gauge and proved the vanishing of the cosmological constant and the non-renormalization theorem [24, 4]. They also computed the four-particle amplitude in another

forthcoming paper [25]. Although the final results are exactly the expected, their computation is quite difficult to follow because of the use of theta functions.¹ Also modular invariance is absurd in their computations because of the complicated dependence between the 2 insertion points (the insertion points are also spin structure dependent).

In the old works [5, 6, 7, 8] on two-loop superstrings, one of the author (with Iengo) used the hyperelliptic representation to do the explicit computation at two loops which is quite explicit and modular invariance is manifest at every stage of the computations. So it is natural to do computations in this language by using the newly established result. In a previous paper [27], we reported the main results of our computation of two loop superstring theory by using hyperelliptic language. In [28] the details for the verification of the vanishing of the cosmological constant and the non-renormalization theorem is given. As we have shown in [28], except one kind term involving the stress energy tensor of the ψ field, all the rest relevant terms containing 3 or less particles are vanishing point-wise in moduli space after summation over spin structures. This non-vanishing term also gives a vanishing contribution to the 3-particle amplitude because of the antisymmetric property of the relevant kinematic factor. As we will see in this paper, the non-renormalization theorem greatly simplifies the computation of the possibly non-vanishing four-particle amplitude.

In this paper we will present the details for the explicit computation of the four-particle amplitude. As announced in [27], we obtained a quite simple and explicit expression for the chiral integrand. For type II superstring theories, we also give an explicit expression for the amplitude. By using this result, we show that the perturbative correction to the R^4 term in type II superstring theories is vanishing at two loops, confirming the computation of Iengo [10] and the conjecture of Green and Gutperle [12] (see also [13, 14]). We leave the proof of the equivalence between the new result and the old result of the four-particle amplitude as a problem of the future. This paper is organized as follows:

In the next section we will recall the relevant results of hyperelliptic representation of the genus 2 Riemann surface and set our notations for all the correlators. In section 3 we recall the results of D'Hoker and Phong for the chiral measure. In section 4 we computed explicitly all the relevant quanti-

¹In [26], the two-loop 4-particle amplitude was also computed by using theta functions. Its relation with the previous explicit result [8] has not been clarified.

ties appearing in the chiral measure and write the chiral integrand (before the summation over spin structures) by using hyperelliptic language. In section 5 we present the computation of the connected terms from the matter supercurrent and show that it gives vanishing contribution to the amplitude. In section 6, we will use the result of [28] and computed the terms with a single $S_n(q)$ factor. We show that the resulting expression is independent of the three arbitrary points $p_{1,2,3}$. In section 7 we compute the rest terms and found an almost total cancellation between the various terms. In section 8, we present the final form for the 4-particle amplitude by combining all the previous results. The final result for the 4-particle amplitude is quite simple. We also show that the result is independent of the insertion points $q_{1,2}$. By using this result, we show that the perturbative correction to the R^4 term in type II superstring theories is vanishing at two loops, confirming the explicit computation of Iengo [10] and the conjecture of Green and Gutperle [12]. Some technical parts are relegated to the appendix. In Appendix A, we collect all the formulas for the $\langle X(z)X(w) \rangle$ correlators. In Appendix B, we present the proof for the summation formulas eq. (78). In Appendix C, we compute explicitly the q dependence of the factor $\mathcal{Z}$. Here we also give some useful formulas which are needed in the transform from the branch point parametrization of the moduli space to the period matrix τ_{ij} .

To conclude this section we note that D'Hoker and Phong have also proved that the cosmological constant and the 1-, 2- and 3-point functions are zero point-wise in moduli space [24]. They have also computed the 4-particle amplitude [25]. The agreement of the results from these two different gauge choices and two different methods of computations would be the final proof of the validity of the new supersymmetric gauge fixing method at two loops.

2 Genus 2 hyperelliptic Riemann surface

First we remind that a genus- g Riemann surface, which is the appropriate world sheet for one and two loops, can be described in full generality by means of the hyperelliptic formalism.² This is based on a representation of the surface as two sheet covering of the complex plane described by the

²Some early works on two loops computation by using hyperelliptic representation are [29, 30, 31, 32, 33, 34, 35] which is by no means the complete list.

equation:

$$y^2(z) = \prod_{i=1}^{2g+2} (z - a_i), \quad (1)$$

The complex numbers a_i , ($i = 1, \dots, 2g + 2$) are the $2g + 2$ branch points, by going around them one passes from one sheet to the other. For two-loop ($g = 2$) three of them represent the moduli of the genus 2 Riemann surface over which the integration is performed, while the other three can be arbitrarily fixed. Another parametrization of the moduli space is given by the period matrix τ . The transformation from the period matrix τ to the branch points can be found in Appendix C.

There are two independent holomorphic differentials on a genus two Riemann surface. In hyperelliptic language these are given explicitly as follows:

$$\Omega_1(z) = \frac{dz}{y(z)}, \quad \Omega_2(z) = \frac{z dz}{y(z)}. \quad (2)$$

These differentials are not normalized in the standard way:

$$\oint_{\alpha_i} \omega_j(z) = \delta_{ij}, \quad \oint_{\beta_i} \omega_j(z) = \tau_{ij}, \quad i, j = 1, 2, \quad (3)$$

where α_i and β_i are the four nontrivial one cycles and τ_{ij} is the 2×2 period matrix. Setting

$$(K)_{ij} = \oint_{\alpha_i} \Omega_j, \quad (4)$$

we have

$$\Omega_i = \omega_j K_{ji}, \quad (5)$$

and

$$\omega_1 = \frac{K_{22}\Omega_1 - K_{21}\Omega_2}{\det K}, \quad (6)$$

$$\omega_2 = \frac{-K_{12}\Omega_1 + K_{11}\Omega_2}{\det K}. \quad (7)$$

At genus 2, by choosing a canonical homology basis of cycles we have the following list of 10 even spin structures:

$$\delta_1 \sim \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \sim (a_1 a_2 a_3 | a_4 a_5 a_6), \quad \delta_2 \sim \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \sim (a_1 a_2 a_4 | a_3 a_5 a_6),$$

$$\begin{aligned}
\delta_3 &\sim \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \sim (a_1 a_2 a_5 | a_3 a_4 a_6), & \delta_4 &\sim \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \sim (a_1 a_2 a_6 | a_3 a_4 a_5), \\
\delta_5 &\sim \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \sim (a_1 a_3 a_4 | a_2 a_5 a_6), & \delta_6 &\sim \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \sim (a_1 a_3 a_5 | a_2 a_4 a_6), \\
\delta_7 &\sim \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \sim (a_1 a_3 a_6 | a_2 a_4 a_5), & \delta_8 &\sim \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \sim (a_1 a_4 a_5 | a_2 a_3 a_6), \\
\delta_9 &\sim \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \sim (a_1 a_4 a_6 | a_2 a_3 a_5), & \delta_{10} &\sim \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \sim (a_1 a_5 a_6 | a_2 a_3 a_4).
\end{aligned}$$

We will denote an even spin structure as $(A_1 A_2 A_3 | B_1 B_2 B_3)$. By convention $A_1 = a_1$. For each even spin structure we have a spin structure dependent factor from determinants which is given as follows [5]:

$$Q_\delta = \prod_{i < j} (A_i - A_j)(B_i - B_j). \quad (8)$$

This is a degree 6 homogeneous polynomials in a_i .

At two loops there are two odd supermoduli and this gives two insertions of supercurrent at two different points x_1 and x_2 . Previously the chiral measure was derived in [36, 18] by a simple projection from the supermoduli space to the even moduli space. This projection doesn't preserve supersymmetry and there is a residual dependence on the two insertion points. This formalism was used in [5, 6, 7, 8]. In these papers we found that it is quite convenient to choose these two insertion points as the two zeroes of a holomorphic abelian differential which are moduli independent points on the Riemann surface. In hyperelliptic language the holomorphic abelian differential can be written generally as follows:

$$\omega_x(z) = \frac{c(z-x) dz}{y(z)}. \quad (9)$$

In hyperelliptic language the two zero points are the same points on the upper and lower sheet of the surface. We denote these two points as $x_1 = x+$ (on the upper sheet) and $x_2 = x-$ (on the lower sheet).

In the following we will give some formulas in hyperelliptic representation which will be used later. First all the relevant correlators are given by³

$$\begin{aligned}
\langle \psi^\mu(z) \psi^\nu(w) \rangle &= -\delta^{\mu\nu} G_{1/2}[\delta](z, w) = -\delta^{\mu\nu} S_\delta(z, w), \\
\langle b(z) c(w) \rangle &= +G_2(z, w), \\
\langle \beta(z) \gamma(w) \rangle &= -G_{3/2}[\delta](z, w),
\end{aligned} \quad (10)$$

³We follow closely the notation of [21].

where

$$S_\delta(z, w) = \frac{1}{z - w} \frac{u(z) + u(w)}{2\sqrt{u(z)u(w)}}, \quad (11)$$

$$u(z) = \prod_{i=1}^3 \left(\frac{z - A_i}{z - B_i} \right)^{1/2}, \quad (12)$$

$$G_2(z, w) = -H(w, z) + \sum_{a=1}^3 H(w, p_a) \varpi_a(z, z), \quad (13)$$

$$H(w, z) = \frac{1}{2(w - z)} \left(1 + \frac{y(w)}{y(z)} \right) \frac{y(w)}{y(z)}, \quad (14)$$

$$G_{3/2}[\delta](z, w) = -P(w, z) + P(w, q_1)\psi_1^*(z) + P(w, q_2)\psi_2^*(z), \quad (15)$$

$$P(w, z) = \frac{1}{\Omega(w)} S_\delta(w, z)\Omega(z), \quad (16)$$

where $\Omega(z)$ is a holomorphic abelian differential satisfying $\Omega(q_{1,2}) \neq 0$ and otherwise arbitrary. These correlators were adapted from [37]. $\varpi_a(z, w)$ are defined in [20] and $\psi_{1,2}^*(z)$ are the two holomorphic $\frac{3}{2}$ -differentials. When no confusion is possible, the dependence on the spin structure $[\delta]$ will not be exhibited. The formulas for the $\langle X(z)X(w) \rangle$ and related correlators are given in Appendix A.

In order to take the limit of $x_{1,2} \rightarrow q_{1,2}$ we need the following expansions:

$$G_{3/2}(x_2, x_1) = \frac{1}{x_1 - q_1} \psi_1^*(x_2) - \psi_1^*(x_2) f_{3/2}^{(1)}(x_2) + O(x_1 - q_1), \quad (17)$$

$$G_{3/2}(x_1, x_2) = \frac{1}{x_2 - q_2} \psi_2^*(x_1) - \psi_2^*(x_1) f_{3/2}^{(2)}(x_1) + O(x_2 - q_2), \quad (18)$$

for $x_{1,2} \rightarrow q_{1,2}$. By using the explicit expression of $G_{3/2}$ in (15) we have

$$f_{3/2}^{(1)}(q_2) = -\frac{\partial_{q_2} S(q_1, q_2)}{S(q_1, q_2)} + \partial \psi_2^*(q_2), \quad (19)$$

$$f_{3/2}^{(2)}(q_1) = \frac{\partial_{q_1} S(q_2, q_1)}{S(q_1, q_2)} + \partial \psi_1^*(q_1) = f_{3/2}^{(1)}(q_2)|_{q_1 \leftrightarrow q_2}. \quad (20)$$

The quantity $\psi_\alpha^*(z)$'s are holomorphic $\frac{3}{2}$ -differentials and are constructed as follows:

$$\psi_\alpha^*(z) = (z - q_\alpha) S(z, q_\alpha) \frac{y(q_\alpha)}{y(z)}, \quad \alpha = 1, 2. \quad (21)$$

For $z = q_{1,2}$ we have

$$\psi_\alpha^*(q_\beta) = \delta_{\alpha\beta}, \quad (22)$$

$$\partial\psi_1^*(q_2) = -\partial\psi_2^*(q_1) = S(q_1, q_2) = \frac{i}{4}S_1(q), \quad (23)$$

$$\partial\psi_1^*(q_1) = \partial\psi_2^*(q_2) = -\frac{1}{2}\Delta_1(q), \quad (24)$$

$$\partial^2\psi_1^*(q_1) = \partial^2\psi_2^*(q_2) = \frac{1}{16}S_1^2(q) + \frac{1}{4}\Delta_1^2(q) + \frac{1}{2}\Delta_2(q), \quad (25)$$

where

$$\Delta_n(x) \equiv \sum_{i=1}^6 \frac{1}{(x - a_i)^n}, \quad (26)$$

$$S_n(x) \equiv \sum_{i=1}^3 \left[\frac{1}{(x - A_i)^n} - \frac{1}{(x - B_i)^n} \right], \quad (27)$$

for $n = 1, 2$. This shows that $\partial\psi_\alpha^*(q_{\alpha+1})$ and $\partial^2\psi_\alpha^*(q_\alpha)$ are spin structure dependent.

The other quantities introduced in [18] which will be used later are as follows:

$$\varpi_1(q_1, q_2) = -\frac{y^2(p_1)}{y^2(q)} \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)}, \quad (28)$$

$$\varpi_2(q_1, q_2) = -\frac{y^2(p_2)}{y^2(q)} \frac{(q - p_1)(q - p_3)}{(p_2 - p_1)(p_2 - p_3)}, \quad (29)$$

$$\varpi_3(q_1, q_2) = -\frac{y^2(p_3)}{y^2(q)} \frac{(q - p_1)(q - p_2)}{(p_3 - p_1)(p_3 - p_2)}, \quad (30)$$

and

$$\begin{aligned} \varpi_1^*(u) &= \frac{y(p_1)}{y(u)} \frac{(up_1 - \frac{1}{2}(u + p_1)(p_2 + p_3) + p_2p_3)}{(p_1 - p_2)(p_1 - p_3)} \\ &= \frac{y(p_1)}{y(u)} \left[1 + \frac{1}{2}(u - p_1) \left(\frac{1}{p_1 - p_2} + \frac{1}{p_1 - p_3} \right) \right], \end{aligned} \quad (31)$$

$$\varpi_2^*(u) = \frac{y(p_2)}{y(u)} \frac{(up_2 - \frac{1}{2}(u + p_2)(p_3 + p_1) + p_1p_3)}{(p_2 - p_3)(p_2 - p_1)}, \quad (32)$$

$$\varpi_3^*(u) = \frac{y(p_3)}{y(u)} \frac{(up_3 - \frac{1}{2}(u + p_3)(p_1 + p_2) + p_1p_2)}{(p_3 - p_1)(p_3 - p_2)}. \quad (33)$$

We note here that $\varpi_1(q_1, q_2) = -1$ and $\varpi_{2,3}(q_1, q_2) = 0$ in the limit $p_1 \rightarrow q_{1,2}$.

3 The chiral measure: the result of D'Hoker and Phong

In this section we will briefly recall the result of D'Hoker and Phong. Previously the chiral measure was derived in [36, 18] by a simple projection from the supermoduli space to the even moduli space. This projection doesn't preserve supersymmetry and there is a residual dependence on the two insertion points. In the new formalism of D'Hoker and Phong, they use a supersymmetric projection to the super-period matrix. For detailed derivation of the chiral measure we refer the reader to their original papers [20, 21, 22, 23]. After making the choice $x_\alpha = q_\alpha$ ($\alpha = 1, 2$), the chiral measure obtained in these papers is:

$$\begin{aligned} \mathcal{A}[\delta] &= i\mathcal{Z} \left\{ 1 + \mathcal{X}_1 + \mathcal{X}_2 + \mathcal{X}_3 + \mathcal{X}_4 + \mathcal{X}_5 + \mathcal{X}_6 \right\}, \\ \mathcal{Z} &= \frac{\langle \Pi_a b(p_a) \Pi_\alpha \delta(\beta(q_\alpha)) \rangle}{\det \omega_I \omega_J(p_a)}, \end{aligned} \quad (34)$$

and the $\mathcal{X}_i$ are given by:

$$\begin{aligned} \mathcal{X}_1 + \mathcal{X}_6 &= \frac{\zeta^1 \zeta^2}{16\pi^2} \left[-\langle \psi(q_1) \cdot \partial X(q_1) \psi(q_2) \cdot \partial X(q_2) \rangle \right. \\ &\quad - \partial_{q_1} G_2(q_1, q_2) \partial \psi_1^*(q_2) + \partial_{q_2} G_2(q_2, q_1) \partial \psi_2^*(q_1) \\ &\quad \left. + 2G_2(q_1, q_2) \partial \psi_1^*(q_2) f_{3/2}^{(1)}(q_2) - 2G_2(q_2, q_1) \partial \psi_2^*(q_1) f_{3/2}^{(2)}(q_1) \right], \end{aligned} \quad (35)$$

$$\begin{aligned} \mathcal{X}_2 + \mathcal{X}_3 &= \frac{\zeta^1 \zeta^2}{8\pi^2} S_\delta(q_1, q_2) \\ &\quad \times \sum_{a=1}^3 \tilde{\omega}_a(q_1, q_2) \left[\langle T(\tilde{p}_a) \rangle + \tilde{B}_2(\tilde{p}_a) + \tilde{B}_{3/2}(\tilde{p}_a) \right], \end{aligned} \quad (36)$$

$$\begin{aligned} \mathcal{X}_4 &= \frac{\zeta^1 \zeta^2}{8\pi^2} S_\delta(q_1, q_2) \sum_{a=1}^3 \left[\partial_{p_a} \partial_{q_1} \ln E(p_a, q_1) \varpi_a^*(q_2) \right. \\ &\quad \left. + \partial_{p_a} \partial_{q_2} \ln E(p_a, q_2) \varpi_a^*(q_1) \right], \end{aligned} \quad (37)$$

$$\begin{aligned} \mathcal{X}_5 &= \frac{\zeta^1 \zeta^2}{16\pi^2} \sum_{a=1}^3 \left[S_\delta(p_a, q_1) \partial_{p_a} S_\delta(p_a, q_2) \right. \\ &\quad \left. - S_\delta(p_a, q_2) \partial_{p_a} S_\delta(p_a, q_1) \right] \varpi_a(q_1, q_2). \end{aligned} \quad (38)$$

Furthermore, $\tilde{B}_2$ and $\tilde{B}_{3/2}$ are given by

$$\tilde{B}_2(w) = -2 \sum_{a=1}^3 \partial_{p_a} \partial_w \ln E(p_a, w) \varpi_a^*(w), \quad (39)$$

$$\tilde{B}_{3/2}(w) = \sum_{\alpha=1}^2 \left(G_2(w, q_\alpha) \partial_{q_\alpha} \psi_\alpha^*(q_\alpha) + \frac{3}{2} \partial_{q_\alpha} G_2(w, q_\alpha) \psi_\alpha^*(q_\alpha) \right). \quad (40)$$

In comparing with [22] we have written $\mathcal{X}_2, \mathcal{X}_3$ together and we didn't split $T(w)$ into different contributions. We also note that in eq. (36) the three arbitrary points $\tilde{p}_a$ ($a = 1, 2, 3$) can be different from the three insertion points p_a 's of the b ghost field. The symbol $\tilde{\varpi}_a$ is obtained from ϖ_a by changing p_a 's to $\tilde{p}_a$'s. In the following computation we will take the limit of $\tilde{p}_1 \rightarrow q_1$. In this limit we have $\tilde{\varpi}_{2,3}(q_1, q_2) = 0$ and $\tilde{\varpi}_1(q_1, q_2) = -1$ as we noted at the end of last section. This choice greatly simplifies the formulas and also make the summation over spin structure doable (see below and [28]).

4 The chiral measure in hyperelliptic language

In this section we will compute explicitly the chiral measure in hyperelliptic language. Here we will not only compute the spin structure dependent parts, but also compute explicitly all the quantities appearing in the chiral measure. In order to do this we will write the chiral measure in hyperelliptic language and then take the limit of $\tilde{p}_1 \rightarrow q_1$.

Let's first start with $\mathcal{X}_5$. We have

$$S(z, q_1) \partial_z S(z, q_2) - S(z, q_2) \partial_z S(z, q_1) = \frac{i}{4(z - q)^2} S_1(z). \quad (41)$$

The exact phase of the above expression is determined by taking the limit $z \rightarrow q_1$. So the chiral integrand of the four-particle amplitude (before the summation over spin structures) from $\mathcal{X}_5$ is⁴

$$\mathcal{A}_5 = \frac{i}{4} \sum_{a=1}^3 \frac{\varpi_a(q_1, q_2)}{(q - p_a)^2} S_1(p_a) \langle \prod_{i=1}^4 V_i(k_i, \epsilon_i; z_i, \bar{z}_i) \rangle, \quad (42)$$

after including the vertex operators.

⁴An overall factor $\frac{\zeta^1 \zeta^2}{16\pi^2}$ was omitted.

For $\mathcal{X}_4$, we need to combine it with the contribution of $\tilde{B}_2(q_1)$ in $\mathcal{X}_2 + \mathcal{X}_3$ in order to get a simpler expression. We have

$$\begin{aligned}
\mathcal{A}_4 + \tilde{B}_2(q_1) \text{ in } \mathcal{X}_2 + \mathcal{X}_3 &= 2 \sum_{a=1}^3 [\partial_{p_a} \partial_{q_1} \ln E(p_a, q_1) + \partial_{p_a} \partial_{q_2} \ln E(p_a, q_2)] \\
&\quad \times \varpi_a^*(q_1) S_1(q_1, q_2) \langle \prod_{i=1}^4 V_i(k_i, \epsilon_i; z_i, \bar{z}_i) \rangle \\
&= \sum_{a=1}^3 \frac{2}{(q - p_a)^2} \varpi_a^*(q_1) S_1(q_1, q_2) \langle \prod_{i=1}^4 V_i(k_i, \epsilon_i; z_i, \bar{z}_i) \rangle, \tag{43}
\end{aligned}$$

where we have used eq. (123) in Appendix A and the fact $\varpi^*(q_2) = -\varpi^*(q_1)$.

For the rest terms in $\mathcal{X}_2 + \mathcal{X}_3$, we first compute the various contributions from the different fields in the stress energy tensor. The total stress energy tensor is:

$$\begin{aligned}
T(z) &= -\frac{1}{2} : \partial_z X(z) \cdot \partial_z X(z) : + \frac{1}{2} : \psi(z) \cdot \partial_z \psi(z) : \\
&\quad - : (\partial b c + 2b \partial c + \frac{1}{2} \partial \beta \gamma + \frac{3}{2} \beta \partial \gamma)(z) : \\
&\equiv T_X(z) + T_\psi(z) + T_{bc}(z) + T_{\beta\gamma}(z), \tag{44}
\end{aligned}$$

in an obvious notations. The various contributions are (following the notations of [21]):

$$T_X(w) = -10T_1(w), \tag{45}$$

$$T_\psi(w) = 5\tilde{g}_{1/2}(w) = \frac{5}{32} (S_1(w))^2, \tag{46}$$

$$T_{bc}(w) = \tilde{g}_2(w) - 2\partial_w f_2(w), \tag{47}$$

$$T_{\beta\gamma}(w) = -\tilde{g}_{3/2}(w) + \frac{3}{2} \partial_w f_{3/2}(w), \tag{48}$$

where

$$f_2(w) = -\frac{3}{4} \Delta_1(w) + \sum_{a=1}^3 H(w, p_a) \varpi_a(w, w), \tag{49}$$

$$\begin{aligned}
\tilde{g}_2(w) &= \frac{5}{16} \Delta_1^2(w) + \frac{3}{8} \Delta_2(w) \\
&\quad + \sum_{a=1}^3 H(w, p_a) \varpi_a(w, w) \left(\frac{1}{w - p_{a+1}} + \frac{1}{w - p_{a+2}} - \Delta_1(w) \right), \tag{50}
\end{aligned}$$

$$f_{3/2}(w) = \frac{\Omega'(w)}{\Omega(w)} + \frac{\Omega(q_1)}{\Omega(w)} S(w, q_1) \psi_1^*(w) + \frac{\Omega(q_2)}{\Omega(w)} S(w, q_2) \psi_2^*(w), \quad (51)$$

$$\begin{aligned} \tilde{g}_{3/2}(w) &= \frac{1}{2} \frac{\Omega''(w)}{\Omega(w)} + \frac{1}{32} (S_1(w))^2 \\ &\quad + \frac{\Omega(q_1)}{\Omega(w)} S(w, q_1) \partial \psi_1^*(w) + \frac{\Omega(q_2)}{\Omega(w)} S(w, q_2) \partial \psi_2^*(w). \end{aligned} \quad (52)$$

As we said in the last section we will take the limit of $w \rightarrow q_1$. In this limit $T_{\beta\gamma}(w)$ is singular and we have the following expansion:

$$T_{\beta\gamma}(w) = -\frac{3/2}{(w - q_1)^2} - \frac{\partial \psi_1^*(q_1)}{w - q_1} - \frac{1}{8} \Delta_1^2(q) - \frac{1}{32} S_1^2(q) + O(w - q_1). \quad (53)$$

The explicit dependence of $G_{3/2}$ on the abelian differential $\Omega(z)$ drops out in the above expression. These singular terms are cancelled by similar singular terms in $\tilde{B}_{3/2}(w)$. By explicit computation we have:

$$\begin{aligned} \tilde{B}_{3/2}(w) &= \frac{3/2}{(w - q_1)^2} + \frac{\partial \psi_1^*(q_1)}{w - q_1} - \frac{1}{4} \Delta_1^2(q) + \frac{3}{4} \Delta_2(q) \\ &\quad - \left(\frac{1}{p_1 - q} \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)} \Delta_1(q) + \dots \right) \\ &\quad - \frac{3}{2} \left(\frac{1}{(p_1 - q)^2} \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)} + \dots \right) + O(w - q_1). \end{aligned} \quad (54)$$

where ... indicates two other terms obtained by cyclicly permutating (p_1, p_2, p_3) . By using the above explicit result we see that the combined contributions of $T_{\beta\gamma}(w)$ and $\tilde{B}_{3/2}(w)$ are non-singular in the limit of $w \rightarrow q_1$. We can then take $\tilde{p}_1 \rightarrow q_1$ in $\mathcal{X}_2 + \mathcal{X}_3$. In this limit only $a = 1$ contributes to $\mathcal{X}_2 + \mathcal{X}_3$. This is because $\tilde{\omega}_{2,3}(q_1, q_2) = 0$ and $\tilde{\omega}_1(q_1, q_2) = -1$. $T_{bc}(w)$ is regular in this limit and it is spin structure independent. It is given as follows:

$$\begin{aligned} T_{bc}(q_1) &= \frac{5}{16} \Delta_1^2(q_1) - \frac{9}{8} \Delta_2(q_1) + \sum_{a=1}^3 \frac{\varpi_a^*(q_1)}{(q_1 - p_a)^2} \\ &\quad + \left\{ \frac{1}{(q_1 - p_1)^2} \left[1 - \frac{1}{2} (q_1 - p_1) \left(\frac{1}{q_1 - p_2} + \frac{1}{q_1 - p_3} + \Delta_1(q_1) \right) \right] \right. \\ &\quad \left. \times \frac{(q_1 - p_2)(q_1 - p_3)}{(p_1 - p_2)(p_1 - p_3)} + \dots \right\}. \end{aligned} \quad (55)$$

where “...” again indicates two other terms obtained by cyclicly permutating (p_1, p_2, p_3) . Combining all the above results together, we have

$$\begin{aligned}
\mathcal{A}_2 + \mathcal{A}_3 = & -2S(q_1, q_2) \left\{ \langle (T_X(q_1) + T_\psi(q_1)) \prod_{i=1}^4 V_i \rangle_c \right. \\
& + \left[\frac{1}{8} (S_1(q))^2 - \frac{1}{16} \Delta_1^2(q) - \frac{3}{8} \Delta_2(q) + \sum_{a=1}^3 \frac{\varpi_a^*(q_1)}{(q-p_a)^2} \right. \\
& - \frac{1}{2} \left\{ \frac{1}{(q-p_1)^2} \left[1 + (q-p_1) \left(\frac{1}{q-p_2} + \frac{1}{q-p_2} - \Delta_1(q_1) \right) \right] \right. \\
& \left. \left. \times \frac{(q-p_2)(q-p_3)}{(p_1-p_2)(p_1-p_3)} + \dots \right\} \right] \langle \prod_{i=1}^4 V_i \rangle \left. \right\}, \tag{56}
\end{aligned}$$

by omitting the contribution of $\tilde{B}_2(q_1)$ which has been included in eq. (43). The subscript “c” indicates that the disconnected contraction of T_ψ with V_i should be omitted. Also by comparing eq. (43) with the above expression we see that it exactly cancels the term with $\varpi_a^*(q_1)$. Writing these expression together we have:

$$\begin{aligned}
\sum_{i=2}^4 \mathcal{A}_i = & -2S(q_1, q_2) \left\{ \langle (T_X(q_1) + T_\psi(q_1)) \prod_{i=1}^4 V_i \rangle \right. \\
& + \left[\frac{1}{8} (S_1(q))^2 - \frac{1}{16} \Delta_1^2(q) - \frac{3}{8} \Delta_2(q) \right. \\
& - \frac{1}{2} \left\{ \frac{1}{(q-p_1)^2} \left[1 + (q-p_1) \left(\frac{1}{q-p_2} + \frac{1}{q-p_2} - \Delta_1(q_1) \right) \right] \right. \\
& \left. \left. \times \frac{(q-p_2)(q-p_3)}{(p_1-p_2)(p_1-p_3)} + \dots \right\} \right] \langle \prod_{i=1}^4 V_i \rangle \left. \right\}. \tag{57}
\end{aligned}$$

Finally we compute $\mathcal{X}_1 + \mathcal{X}_6$ explicitly. By using the explicit results given in eqs. (19)–(20), we have:

$$\begin{aligned}
\mathcal{A}_1 + \mathcal{A}_6 = & -\langle \psi(q_1) \cdot \partial X(q_1) \psi(q_2) \cdot \partial X(q_2) \prod_{i=1}^4 V_i \rangle \\
& -(\partial_{q_1} G_2(q_1, q_2) + \partial_{q_2} G_2(q_2, q_1)) S(q_1, q_2) \langle \prod_{i=1}^4 V_i \rangle \\
& + 2(G_2(q_1, q_2) + G_2(q_2, q_1)) \\
& \times (\partial \psi_1^*(q_1) S(q_1, q_2) - \partial_{q_2} S(q_1, q_2)) \langle \prod_{i=1}^4 V_i \rangle. \tag{58}
\end{aligned}$$

Here it is important that the factor $\partial\psi_1^*(q_2)$ cancels the factor $S(q_1, q_2)$ appearing in the denominator of $f_{3/2}^{(1)}(q_2)$. The G_2 's appearing in the above expression can be computed explicitly by using eq. (13). The expressions that we will need are given as follows:

$$G_2(q_1, q_2) + G_2(q_2, q_1) = -\frac{1}{2} \Delta_1(q) + \left[\frac{1}{q-p_1} \frac{(q-p_2)(q-p_3)}{(p_1-p_2)(p_1-p_3)} + \dots \right], \quad (59)$$

$$\begin{aligned} \partial_{q_1} G_2(q_1, q_2) + \partial_{q_2} G_2(q_2, q_1) &= \frac{3}{8} \Delta_1^2(q) + \frac{1}{4} \Delta_2(q) \\ &+ \left[\frac{1}{q-p_1} \left(\frac{1}{q-p_2} + \frac{1}{q-p_3} - \Delta_1(q) \right) \frac{(q-p_2)(q-p_3)}{(p_1-p_2)(p_1-p_3)} + \dots \right]. \end{aligned} \quad (60)$$

Now we turn to the computation of the 4-particle amplitude. The computation was split into three sections.

5 The 4-particle amplitude I: the vanishing of the connected term $\langle \psi(q_1) \psi(q_2) \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle_c$

For graviton and the antisymmetric tensor we use the following vertex operator (left part only):

$$V_i(k_i, \epsilon_i, z_i) = (\epsilon_i \cdot \partial X(z_i) + i k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i)) e^{i k_i \cdot X(z_i, \bar{z}_i)}. \quad (61)$$

Because the vertex operator doesn't contain any ghost fields, all terms involving ghost fields can be explicitly computed which we have done in the above. For the computation of amplitudes of other kinds of particles (like fermions), one either resorts to supersymmetry or can use similar method which was used in [38, 39] to compute the fermionic amplitude.

Before we do explicit computations, we will show that the connected term $\langle \psi(q_1) \psi(q_2) \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle_c$ gives vanishing contributions by doing summation over spin structures. This result was first discovered in [8, 7]. Let's recall the argument here.

First we note that all the connected contractions are the following three kinds:

$$A_\delta = \langle \psi(q_1)\psi(z_1) \rangle \langle \psi(z_1)\psi(q_2) \rangle \\ \times \langle \psi(z_2)\psi(z_3) \rangle \langle \psi(z_3)\psi(z_4) \rangle \langle \psi(z_4)\psi(z_2) \rangle, \quad (62)$$

$$B_\delta = \langle \psi(q_1)\psi(z_1) \rangle \langle \psi(z_1)\psi(z_2) \rangle \langle \psi(z_2)\psi(q_2) \rangle (\langle \psi(z_3)\psi(z_4) \rangle)^2, \quad (63)$$

$$C_\delta = \langle \psi(q_1)\psi(z_1) \rangle \langle \psi(z_1)\psi(z_2) \rangle \\ \times \langle \psi(z_2)\psi(z_3) \rangle \langle \psi(z_3)\psi(z_4) \rangle \langle \psi(z_4)\psi(q_2) \rangle, \quad (64)$$

$$(65)$$

or sometimes the expressions permuted among z_1, z_2, z_3 and z_4 . By using the explicit expression of $\langle \psi(z)\psi(w) \rangle$ and noting $u(q_2) = -u(q_1)$, we have

$$A_\delta \propto 2 \left[\frac{u(q_1)}{u(z_1)} - \frac{u(z_1)}{u(q_1)} \right] + \sum_{i \neq j=2}^4 \left[\frac{u(q_1)u(z_i)}{u(z_1)u(z_j)} - \frac{u(z_1)u(z_j)}{u(q_1)u(z_i)} \right], \quad (66)$$

$$B_\delta \propto 2 \sum_{i=1}^2 \left[\frac{u(q_1)}{u(z_i)} - \frac{u(z_i)}{u(q_1)} \right] + 2 \left[\frac{u(z_1)}{u(z_2)} - \frac{u(z_2)}{u(z_1)} \right] \\ + \sum_{i=1}^2 \sum_{k \neq l=3}^4 \left[\frac{u(q_1)u(z_k)}{u(z_i)u(z_l)} - \frac{u(z_i)u(z_l)}{u(q_1)u(z_k)} \right] \\ + \sum_{k \neq l=3}^4 \left[\frac{u(z_1)u(z_k)}{u(z_2)u(z_l)} - \frac{u(z_2)u(z_l)}{u(z_1)u(z_k)} \right], \quad (67)$$

$$C_\delta \propto \sum_{i=1}^4 \left[\frac{u(q_1)}{u(z_i)} - \frac{u(z_i)}{u(q_1)} \right] + \sum_{i < j=1}^4 \left[\frac{u(z_i)}{u(z_j)} - \frac{u(z_j)}{u(z_i)} \right] \\ + \left[\frac{u(z_1)u(z_3)}{u(z_2)u(z_4)} - \frac{u(z_2)u(z_4)}{u(z_1)u(z_3)} \right] + \sum_{i=3}^4 \left[\frac{u(q_1)u(z_2)}{u(z_1)u(z_i)} - \frac{u(z_1)u(z_i)}{u(q_1)u(z_2)} \right] \\ + \sum_{i=1}^2 \left[\frac{u(q_1)u(z_3)}{u(z_i)u(z_4)} - \frac{u(z_i)u(z_4)}{u(q_1)u(z_3)} \right]. \quad (68)$$

By using the following “vanishing identities”:

$$\sum_{\delta} \eta_{\delta} Q_{\delta} \left[\frac{u(z)}{u(w)} - \frac{u(w)}{u(z)} \right] = 0, \quad (69)$$

$$\sum_{\delta} \eta_{\delta} Q_{\delta} \left[\frac{u(z_1)u(z_2)}{u(z_3)u(z_4)} - \frac{u(z_3)u(z_4)}{u(z_1)u(z_2)} \right] = 0, \quad (70)$$

given in [28], we readily show that

$$\sum_{\delta} \eta_{\delta} Q_{\delta} A_{\delta} = \sum_{\delta} \eta_{\delta} Q_{\delta} B_{\delta} = \sum_{\delta} \eta_{\delta} Q_{\delta} C_{\delta} = 0. \quad (71)$$

This shows that all the connected terms in $\langle \psi(q_1) \psi(q_2) \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle_c$ give vanishing contributions to the 4-particle amplitude after summation over spin structures, and can be neglected. For the first term in eq. (58) we only need to compute the disconnected term $S(q_1, q_2) \langle \partial X(q_1) \cdot \partial X(q_2) \prod_{i=1}^4 V_i \rangle$.

6 The 4-particle amplitude II: the $S_n(q) \langle \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle$ terms

The next step in the explicit calculation of the 4-particle amplitude is to compute the terms with a single $S_n(q)$ (or $S_1(p_a)$ from $\mathcal{A}_5$) in the summation over spin structures, i.e. terms like $S_n(q) \langle \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle$. Other terms with less ψ fields automatically give 0 after summation over spin structures (the non-renormalization theorem verified in [28]).

The first step in this computation is to do the contraction for the ψ fields. We have

$$\begin{aligned} \langle \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle &= K_1(1, 2, 3, 4) (S(z_1, z_2) S(z_3, z_4))^2 \\ &\quad + K_1(1, 3, 4, 2) (S(z_1, z_3) S(z_2, z_4))^2 \\ &\quad + K_1(1, 4, 2, 3) (S(z_1, z_4) S(z_2, z_3))^2 \\ &\quad + K_2(1, 2, 3, 4) S(z_1, z_2, z_3, z_4) \\ &\quad + K_2(1, 3, 4, 2) S(z_1, z_3, z_4, z_2) \\ &\quad + K_2(1, 4, 2, 3) S(z_1, z_4, z_2, z_3), \end{aligned} \quad (72)$$

where

$$S(z_1, z_2, z_3, z_4) = S(z_1, z_2) S(z_2, z_3) S(z_3, z_4) S(z_4, z_1), \quad (73)$$

$$\begin{aligned} K_1(1, 2, 3, 4) &= (k_1 \cdot k_2 \epsilon_1 \cdot \epsilon_2 - k_1 \cdot \epsilon_2 k_2 \cdot \epsilon_1) \\ &\quad \times (k_3 \cdot k_4 \epsilon_3 \cdot \epsilon_4 - k_3 \cdot \epsilon_4 k_4 \cdot \epsilon_3), \end{aligned} \quad (74)$$

$$\begin{aligned} K_2(1, 2, 3, 4) &= -k_1 \cdot k_2 k_3 \cdot k_4 \epsilon_1 \cdot \epsilon_4 \epsilon_2 \cdot \epsilon_3 - k_1 \cdot k_4 k_2 \cdot k_3 \epsilon_1 \cdot \epsilon_2 \epsilon_3 \cdot \epsilon_4 \\ &\quad + k_1 \cdot k_2 (k_3 \cdot \epsilon_4 k_4 \cdot \epsilon_1 \epsilon_2 \cdot \epsilon_3 + k_3 \cdot \epsilon_2 k_4 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_4 - k_3 \cdot \epsilon_2 k_4 \cdot \epsilon_1 \epsilon_3 \cdot \epsilon_4) \end{aligned}$$

$$\begin{aligned}
& +k_1 \cdot k_4(k_2 \cdot \epsilon_1 k_3 \cdot \epsilon_2 \epsilon_3 \cdot \epsilon_4 + k_2 \cdot \epsilon_3 k_3 \cdot \epsilon_4 \epsilon_1 \cdot \epsilon_2 - k_2 \cdot \epsilon_1 k_3 \cdot \epsilon_4 \epsilon_2 \cdot \epsilon_3) \\
& +k_2 \cdot k_3(k_1 \cdot \epsilon_2 k_4 \cdot \epsilon_1 \epsilon_3 \cdot \epsilon_4 + k_1 \cdot \epsilon_4 k_4 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_2 - k_1 \cdot \epsilon_2 k_4 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_4) \\
& +k_3 \cdot k_4(k_1 \cdot \epsilon_2 k_2 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_4 + k_1 \cdot \epsilon_4 k_2 \cdot \epsilon_1 \epsilon_2 \cdot \epsilon_3 - k_1 \cdot \epsilon_4 k_2 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_2) \\
& -k_1 \cdot \epsilon_2 k_2 \cdot \epsilon_3 k_3 \cdot \epsilon_4 k_4 \cdot \epsilon_1 - k_1 \cdot \epsilon_4 k_2 \cdot \epsilon_1 k_3 \cdot \epsilon_2 k_4 \cdot \epsilon_3.
\end{aligned} \tag{75}$$

We note that the above two kinematic factors have a symmetry under the simultaneous interchange of $1 \leftrightarrow 4$ and $2 \leftrightarrow 3$. K_2 is also invariant under the cyclic permutations of $(1, 2, 3, 4)$, i.e., $K_2(1, 2, 3, 4) = K_2(2, 3, 4, 1) = K_2(3, 4, 1, 2)$, etc.

By using the explicit expression for the Szegő kernel $S(z_1, z_2)$ given in eq. (11), we have

$$\begin{aligned}
(S(z_1, z_2)S(z_3, z_4))^2 &= \frac{1}{16 z_{12}^2 z_{34}^2} \left\{ 4 + 2 \left[\frac{u(z_1)}{u(z_2)} + \frac{u(z_2)}{u(z_1)} \right] + 2 \left[\frac{u(z_3)}{u(z_4)} + \frac{u(z_4)}{u(z_3)} \right] \right. \\
& \quad \left. + \left[\frac{u(z_1)u(z_3)}{u(z_2)u(z_4)} + \frac{u(z_2)u(z_4)}{u(z_1)u(z_3)} \right] + \left[\frac{u(z_1)u(z_4)}{u(z_2)u(z_3)} + \frac{u(z_2)u(z_3)}{u(z_1)u(z_4)} \right] \right\}, \tag{76}
\end{aligned}$$

$$\begin{aligned}
S(z_1, z_2, z_3, z_4) &= \frac{1}{16 z_{12} z_{23} z_{34} z_{41}} \left\{ 2 + \sum_{i < j=1}^4 \left[\frac{u(z_i)}{u(z_j)} + \frac{u(z_j)}{u(z_i)} \right] \right. \\
& \quad \left. + \left[\frac{u(z_1)u(z_3)}{u(z_2)u(z_4)} + \frac{u(z_2)u(z_4)}{u(z_1)u(z_3)} \right] \right\}. \tag{77}
\end{aligned}$$

By using the “vanishing identities” proved in [28], most of the terms are 0 after summation over spin structures. In order to do the summation over spin structure for the rest non-vanishing expression we need the following summation formula:

$$\begin{aligned}
\sum_{\delta} \eta_{\delta} Q_{\delta} \left\{ \frac{u(z_1)u(z_2)}{u(z_3)u(z_4)} - (-1)^n \frac{u(z_1)u(z_2)}{u(z_3)u(z_4)} \right\} (S_m(x))^n \\
= \frac{2P(a) \prod_{i=1}^2 \prod_{j=3}^4 (z_i - z_j) \prod_{i=1}^4 (x - z_i)}{y^2(x) \prod_{i=1}^4 y(z_i)} \times C_{n,m}, \tag{78}
\end{aligned}$$

where

$$C_{1,1} = 1, \tag{79}$$

$$C_{2,1} = -2(\tilde{z}_1 + \tilde{z}_2 - \tilde{z}_3 - \tilde{z}_4), \tag{80}$$

$$C_{1,2} = \Delta_1(x) - \sum_{k=1}^4 \tilde{z}_k, \tag{81}$$

$$C_{3,1} = 2\Delta_2(x) - \Delta_1^2(x) + 2\Delta_1(x) \sum_{k=1}^4 \tilde{z}_k + 4(\tilde{z}_1\tilde{z}_2 - 2(\tilde{z}_1 + \tilde{z}_2)(\tilde{z}_3 + \tilde{z}_4) + \tilde{z}_3\tilde{z}_4), \quad (82)$$

$$\tilde{z}_k = \frac{1}{x - z_k}, \quad (83)$$

$$P(a) = \prod_{i < j} (a_i - a_j). \quad (84)$$

$C_{1,1}$ and $C_{1,2}$ were firstly derived in [8] (see also [7]). Although other values of n, m also gives modular invariant expressions, the results are quite complex.⁵ Fortunately we only need to use the above listed results. The proof of the above summation formula can be found in Appendix B.

By using these result one easily prove the following:

$$\begin{aligned} \sum_{\delta} \eta_{\delta} Q_{\delta} S_n(x) S(z_1, z_2, z_3, z_4) &= \sum_{\delta} \eta_{\delta} Q_{\delta} S_n(x) (S(z_1, z_2) S(z_3, z_4))^2 \\ &= \frac{P(a)}{8y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} \begin{cases} 1, & n = 1, \\ \Delta_1(x) - \sum_{i=1}^4 \tilde{z}_i, & n = 2. \end{cases} \end{aligned} \quad (85)$$

By using this result and eq. (72), we have

$$\sum_{\delta} \eta_{\delta} Q_{\delta} S_n(x) \langle \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle \quad (86)$$

$$= K(k_i, \epsilon_i) \frac{P(a)}{8y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} \times \begin{cases} 1, & n = 1, \\ \Delta_1(x) - \sum_{i=1}^4 \tilde{z}_i, & n = 2, \end{cases} \quad (87)$$

where

$$\begin{aligned} K(k_i, \epsilon_i) &\equiv K_1(1, 2, 3, 4) + K_1(1, 3, 4, 2) + K_1(1, 4, 2, 3) \\ &\quad + K_2(1, 2, 3, 4) + K_2(1, 3, 4, 2) + K_2(1, 4, 2, 3) \\ &= -\frac{1}{4}(s t \epsilon_1 \cdot \epsilon_3 \epsilon_2 \cdot \epsilon_4 + s u \epsilon_2 \cdot \epsilon_3 \epsilon_1 \cdot \epsilon_4 + t u \epsilon_1 \cdot \epsilon_2 \epsilon_3 \cdot \epsilon_4) \\ &\quad + \frac{1}{2} s (\epsilon_1 \cdot k_4 \epsilon_3 \cdot k_2 \epsilon_2 \cdot \epsilon_4 + \epsilon_2 \cdot k_3 \epsilon_4 \cdot k_1 \epsilon_1 \cdot \epsilon_3 \\ &\quad + \epsilon_1 \cdot k_3 \epsilon_4 \cdot k_2 \epsilon_2 \cdot \epsilon_3 + \epsilon_2 \cdot k_4 \epsilon_3 \cdot k_1 \epsilon_1 \cdot \epsilon_4) \end{aligned}$$

⁵This is due to the non-vanishing of the summation over spin structures when we set $z_1 = z_3$ or $z_1 = z_4$, etc.

$$\begin{aligned}
& + \frac{1}{2} t (\epsilon_2 \cdot k_1 \epsilon_4 \cdot k_3 \epsilon_1 \cdot \epsilon_3 + \epsilon_3 \cdot k_4 \epsilon_1 \cdot k_2 \epsilon_2 \cdot \epsilon_4 \\
& \quad + \epsilon_2 \cdot k_4 \epsilon_1 \cdot k_3 \epsilon_3 \cdot \epsilon_4 + \epsilon_3 \cdot k_1 \epsilon_4 \cdot k_2 \epsilon_1 \cdot \epsilon_2) \\
& + \frac{1}{2} u (\epsilon_1 \cdot k_2 \epsilon_4 \cdot k_3 \epsilon_2 \cdot \epsilon_3 + \epsilon_3 \cdot k_4 \epsilon_2 \cdot k_1 \epsilon_1 \cdot \epsilon_4 \\
& \quad + \epsilon_1 \cdot k_4 \epsilon_2 \cdot k_3 \epsilon_1 \cdot \epsilon_4 + \epsilon_3 \cdot k_2 \epsilon_4 \cdot k_1 \epsilon_1 \cdot \epsilon_2), \tag{88}
\end{aligned}$$

is the standard kinematic factor (left part only) appearing at tree and one loop computation of superstring theories [1, 8, 7]. So apart from an overall factor and the kinematic factor, each appearance of $S_n(x) \langle \prod_{i=1}^4 V_i(k_i, \epsilon; z_i, \bar{z}_i) \rangle$ in the chiral integrand $\mathcal{A}_i$ can be substituted by either 1 or $\Delta_1(x) - \sum_{i=1}^4 \tilde{z}_i$ for $n = 1$ or 2. After this has been done we have the following form for the chiral integrand:⁶

$$\begin{aligned}
\mathcal{A} &= \sum_{a=1}^3 \frac{\varpi_a(q_1, q_2)}{(q - p_a)^2} \times \frac{y^2(q)}{y(p_a)} \prod_{i=1}^4 \frac{p_a - z_i}{q - z_i} \langle \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \\
&+ \langle (\partial X(q_1) \cdot (\partial X(q_1) + \partial X(q_2))) \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \\
&+ \left\{ \frac{1}{8} \Delta_1^2(q) + \frac{3}{4} \Delta_2(q) + \left\{ \left[\frac{1}{(q - p_1)^2} + \frac{1}{q - p_1} \right. \right. \right. \\
&\quad \times \left. \left. \left(\frac{1}{q - p_2} + \frac{1}{q - p_3} - \Delta_1(q) \right) \right] \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)} + \dots \right\} \\
&- \frac{3}{8} \Delta_1^2(q) - \frac{1}{4} \Delta_2(q) - \left[\frac{1}{q - p_1} \left(\frac{1}{q - p_2} + \frac{1}{q - p_3} - \Delta_1(q) \right) \right. \\
&\quad \times \left. \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)} + \dots \right] + \frac{1}{2} \Delta_1(q) \sum_{k=1}^4 \tilde{z}_k \\
&- \left. \left[\frac{1}{q - p_1} \frac{(q - p_2)(q - p_3)}{(p_1 - p_2)(p_1 - p_3)} + \dots \right] \sum_{k=1}^4 \tilde{z}_k \right\} \langle \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \\
&= \langle : \partial X(q_1) \cdot (\partial X(q_1) + \partial X(q_2)) : \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle
\end{aligned}$$

⁶omitting the $(S_1(q))^3$ and T_ψ terms and an overall factor $\frac{P(a)}{8y^2(q)} \prod_{i=1}^4 \frac{q - z_i}{y(z_i)}$. So the contribution from $\mathcal{A}_5$ has an extra factor of $\frac{y^2(q)}{y^2(p_a)} \prod_{i=1}^4 \frac{p_a - z_i}{q - z_i}$. $\tilde{z}_k$ in the following denotes $\frac{1}{q - z_k}$.

$$+ \left[\frac{1}{2} \Delta_2(q) - \frac{1}{4} \Delta_1^2(q) + \frac{1}{2} \Delta_1(q) \sum_{i=1}^4 \tilde{z}_k - \sum_{k < l} \tilde{z}_k \tilde{z}_l \right] \left\langle \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \right\rangle, \quad (89)$$

and the complete chiral integrand is

$$\begin{aligned} \mathcal{A} = & K(k_i, \epsilon_i) \left\{ \langle : \partial X(q_1) \cdot (\partial X(q_1) + \partial X(q_2)) : \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \right. \\ & + \left[\frac{1}{2} \Delta_2(q) - \frac{1}{4} \Delta_1^2(q) + \frac{1}{2} \Delta_1(q) \sum_{i=1}^4 \tilde{z}_k - \sum_{k < l} \tilde{z}_k \tilde{z}_l \right] \\ & \times \left\langle \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \right\rangle \left. \right\} \times \frac{P(a)}{8 y^2(q)} \prod_{i=1}^4 \frac{q - z_i}{y(z_i)} \\ & - \left\langle \left\langle \frac{1}{4} (S_1(q))^3 \left\langle \prod_{i=1}^4 V_i \right\rangle + S_1(q) \langle : \psi(q_1) \cdot \partial \psi(q_1) : \prod_{i=1}^4 V_i \rangle_c \right\rangle \right\rangle_s. \quad (90) \end{aligned}$$

Here we have reinserted the overall factor. The subscript c indicates that the self contraction of $\psi(q_1)$ with $\partial \psi(q_1)$ is omitted and $\langle \langle \dots \rangle \rangle_s$ indicates that appropriate summation over spin structure should be carried out. As one can see from the above results, all the dependence on p_a 's drops out. This is a very strong check for the validity of the new supersymmetric gauge fixing method at two loops.

7 The 4-particle amplitude III: terms from $(S_1(q))^3$ and T_ψ

To compute the last term in eq. (90), we need the following summation formulas:

$$\begin{aligned} & \sum_{\delta} \eta_{\delta} Q_{\delta} (S(x, z_1) S(z_1, z_2) \partial_x S(z_2, x) + (z_1 \leftrightarrow z_2)) (S(z_3, z_4))^2 S_1(x) \\ & = - \frac{P(a)}{16 y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} (\tilde{z}_{14} \tilde{z}_{23} + \tilde{z}_{13} \tilde{z}_{24}), \quad (91) \end{aligned}$$

$$\begin{aligned} & \sum_{\delta} \eta_{\delta} Q_{\delta} (S(x, z_1) S(z_1, z_2) S(z_2, z_3) S(z_3, z_4) \partial_x S(z_4, x) \\ & + (z_1 \leftrightarrow z_4, z_2 \leftrightarrow z_3)) S_1(x) = \frac{P(a)}{16 y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} \tilde{z}_{14} \tilde{z}_{23}, \quad (92) \end{aligned}$$

$$\begin{aligned}
& \sum_{\delta} \eta_{\delta} Q_{\delta} (S(z_1, z_2) S(z_3, z_4))^2 (S_1(x))^3 \\
&= \frac{P(a)}{8 y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} \left[2\Delta_2(x) - \Delta_1^2(x) + 2\Delta_1(x) \sum_{i=1}^4 \tilde{z}_i \right. \\
&\quad \left. + 4 \sum_{k < l} \tilde{z}_k \tilde{z}_l - 24(\tilde{z}_1 \tilde{z}_2 + \tilde{z}_3 \tilde{z}_4) \right], \tag{93}
\end{aligned}$$

$$\begin{aligned}
& \sum_{\delta} \eta_{\delta} Q_{\delta} S(z_1, z_2) S(z_2, z_3) S(z_3, z_4) S(z_4, z_1) (S_1(x))^3 \\
&= \frac{P(a)}{8 y^2(x)} \prod_{i=1}^4 \frac{x - z_i}{y(z_i)} \left[2\Delta_2(x) - \Delta_1^2(x) + 2\Delta_1(x) \sum_{i=1}^4 \tilde{z}_i \right. \\
&\quad \left. + 4 \sum_{k < l} \tilde{z}_k \tilde{z}_l - 12(\tilde{z}_1 + \tilde{z}_3)(\tilde{z}_2 + \tilde{z}_4) \right]. \tag{94}
\end{aligned}$$

All these formulas can be derived by using the formulas given in eq. (78) and the following useful formula for the derivative of the Szegő kernel:

$$\partial_x S(z, x) = \frac{1}{2(z-x)^2} \frac{u(z) + u(x)}{\sqrt{u(z)u(x)}} - \frac{S_1(x)}{8(z-x)} \frac{u(z) - u(x)}{\sqrt{u(z)u(x)}}. \tag{95}$$

Now we compute the last term explicitly in eq. (90). First we have the following contractions:

$$\begin{aligned}
& \langle \psi(x) \cdot \partial \psi(x) \prod_{i=1}^4 k_i \cdot \psi(z_i) \epsilon_i \cdot \psi(z_i) \rangle \\
&= -2 CS_1(1, 2, 3, 4) K_1(1, 2, 3, 4) - 2 CS_1(1, 3, 4, 2) K_1(1, 3, 4, 2) \\
&\quad - 2 CS_1(1, 3, 4, 2) K_1(1, 3, 4, 2) + CS_2(1, 2, 3, 4) K_2(1, 2, 3, 4) \\
&\quad + CS_2(1, 3, 4, 2) K_2(1, 3, 4, 2) + CS_2(1, 3, 4, 2) K_2(1, 3, 4, 2), \tag{96}
\end{aligned}$$

where

$$\begin{aligned}
CS_1(1, 2, 3, 4) &= (S(x, z_1) S(z_1, z_2) \partial_x S(z_2, x) + (z_1 \leftrightarrow z_2)) (S(z_3, z_4))^2 \\
&\quad + (S(x, z_3) S(z_3, z_4) \partial_x S(z_4, x) + (z_3 \leftrightarrow z_4)) (S(z_1, z_2))^2, \tag{97}
\end{aligned}$$

$$\begin{aligned}
CS_2(1, 2, 3, 4) &= S(x, z_1, z_2, z_3, z_4) + S(x, z_2, z_3, z_4, z_1) \\
&\quad + S(x, z_3, z_4, z_1, z_2) + S(x, z_4, z_1, z_2, z_3) \\
&\quad + S(x, z_3, z_2, z_1, z_4) + S(x, z_4, z_3, z_2, z_1) \\
&\quad + S(x, z_4, z_3, z_2, z_1) + S(x, z_4, z_3, z_2, z_1), \tag{98}
\end{aligned}$$

$$S(x, z_1, z_2, z_3, z_4) \equiv S(x, z_1) S(z_1, z_2) S(z_2, z_3) S(z_3, z_4) \partial_x S(z_4, x), \tag{99}$$

and the kinematic factors $K_{1,2}(1, 2, 3, 4)$ are defined as before. By using eq. (91), one sees that $CS_1(1, 2, 3, 4)$ gives a factor:

$$\begin{aligned} & -\frac{1}{2}(\tilde{z}_{14}\tilde{z}_{23} + \tilde{z}_{13}\tilde{z}_{24} + (\tilde{z}_1 \leftrightarrow \tilde{z}_3, \tilde{z}_2 \leftrightarrow \tilde{z}_4)) \\ & = -3(\tilde{z}_1\tilde{z}_2 + \tilde{z}_3\tilde{z}_4) + \sum_{k<l} \tilde{z}_k\tilde{z}_l, \end{aligned} \quad (100)$$

and by using eq. (92), one sees that $CS_1(1, 2, 3, 4)$ gives a factor:

$$\begin{aligned} & \frac{1}{2}(\tilde{z}_{14}\tilde{z}_{23} + \tilde{z}_{21}\tilde{z}_{34} + \tilde{z}_{32}\tilde{z}_{41} + \tilde{z}_{43}\tilde{z}_{12}) \\ & = 3(\tilde{z}_1 + \tilde{z}_3)(\tilde{z}_2 + \tilde{z}_4) - 2\sum_{k<l} \tilde{z}_k\tilde{z}_l. \end{aligned} \quad (101)$$

By using these results one computes the connected contribution of the T_ψ term as follows:

$$\begin{aligned} \mathcal{A}_{T_\psi} &= - \left[-2(-3(\tilde{z}_1\tilde{z}_2 + \tilde{z}_3\tilde{z}_4) + \sum_{k<l} \tilde{z}_k\tilde{z}_l)K_1(1, 2, 3, 4) \right. \\ & \quad -2(-3(\tilde{z}_1\tilde{z}_3 + \tilde{z}_2\tilde{z}_4) + \sum_{k<l} \tilde{z}_k\tilde{z}_l)K_1(1, 3, 4, 2) \\ & \quad +2(-3(\tilde{z}_1\tilde{z}_4 + \tilde{z}_2\tilde{z}_3) + \sum_{k<l} \tilde{z}_k\tilde{z}_l)K_1(1, 4, 2, 3) \\ & \quad +3(\tilde{z}_1 + \tilde{z}_3)(\tilde{z}_2 + \tilde{z}_4) - 2\sum_{k<l} \tilde{z}_k\tilde{z}_l)K_2(1, 2, 3, 4) \\ & \quad +3(\tilde{z}_1 + \tilde{z}_4)(\tilde{z}_2 + \tilde{z}_3) - 2\sum_{k<l} \tilde{z}_k\tilde{z}_l)K_2(1, 3, 4, 2) \\ & \quad \left. +3(\tilde{z}_1 + \tilde{z}_2)(\tilde{z}_3 + \tilde{z}_4) - 2\sum_{k<l} \tilde{z}_k\tilde{z}_l)K_2(1, 4, 2, 3) \right] \\ &= 2K(k_i, \epsilon_i) \sum_{k<l} \tilde{z}_k\tilde{z}_l \\ & \quad -6(\tilde{z}_1\tilde{z}_2 + \tilde{z}_3\tilde{z}_4)K_1(1, 2, 3, 4) - 6(\tilde{z}_1\tilde{z}_3 + \tilde{z}_2\tilde{z}_4)K_1(1, 3, 4, 2) \\ & \quad -6(\tilde{z}_1\tilde{z}_4 + \tilde{z}_2\tilde{z}_3)K_1(1, 4, 2, 3) - 3(\tilde{z}_1 + \tilde{z}_3)(\tilde{z}_2 + \tilde{z}_4)K_2(1, 2, 3, 4) \\ & \quad -3(\tilde{z}_1 + \tilde{z}_4)(\tilde{z}_2 + \tilde{z}_3)K_2(1, 3, 4, 2) - 3(\tilde{z}_1 + \tilde{z}_2)(\tilde{z}_3 + \tilde{z}_4), \end{aligned} \quad (102)$$

apart from an overall factor.

The $(S_1(q))^3$ contribution can also be computed. We need to use eq. (72) for the ψ contraction and then eqs. (93) and (94) for the summation over

spin structures. The result is

$$\begin{aligned}
\mathcal{A}_{S_1^3} = & -\frac{1}{4} \left[(2\Delta_2(q) - \Delta_1^2(q) + 2\Delta_1(q) \sum_k \tilde{z}_k + 4 \sum_{k<l} \tilde{z}_k \tilde{z}_l) K(k_i, \epsilon_i) \right. \\
& -24(\tilde{z}_1 \tilde{z}_2 + \tilde{z}_3 \tilde{z}_4) K_1(1, 2, 3, 4) - 24(\tilde{z}_1 \tilde{z}_3 + \tilde{z}_2 \tilde{z}_4) K_1(1, 3, 4, 2) \\
& -24(\tilde{z}_1 \tilde{z}_4 + \tilde{z}_2 \tilde{z}_3) K_1(1, 4, 2, 3) \\
& -12(\tilde{z}_1 + \tilde{z}_3)(\tilde{z}_2 + \tilde{z}_4) K_2(1, 2, 3, 4) \\
& -12(\tilde{z}_1 + \tilde{z}_4)(\tilde{z}_2 + \tilde{z}_3) K_2(1, 3, 4, 2) \\
& \left. -12(\tilde{z}_1 + \tilde{z}_2)(\tilde{z}_3 + \tilde{z}_4) K_2(1, 4, 2, 3) \right]. \tag{103}
\end{aligned}$$

By using eq. (102) and eq. (103), we have

$$\begin{aligned}
\mathcal{A}_{T_\psi} + \mathcal{A}_{S_1^3} = & \left[-\frac{1}{2} \Delta_2(q) + \frac{1}{4} \Delta_1^2(q) - \frac{1}{2} \Delta_1(q) \sum_k \tilde{z}_k + \sum_{k<l} \tilde{z}_k \tilde{z}_l \right] \\
& \times K(k_i, \epsilon_i) \langle \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \frac{P(a)}{8y^2(q)} \prod_{i=1}^4 \frac{q - z_i}{y(z_i)}, \tag{104}
\end{aligned}$$

by including the overall factor. The above contribution also has the standard kinematic factor and it exactly cancels the second part of the first term of eq. (90).

By using eq. (90) and eq. (104), we found that final result for the complete chiral integrand of the four-particle amplitude is quite simple and it is given as follows:

$$\mathcal{A} = K(k_i, \epsilon_i) \langle : \partial X(q_1) \cdot (\partial X(q_1) + \partial X(q_2)) : \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \prod_{i=1}^4 \frac{q - z_i}{y(z_i)}, \tag{105}$$

where an overall factor $\frac{P(a)}{8y^2(q)}$ is omitted. This factor is cancelled by an identical factor from the factor $\mathcal{Z}$ (see [8] and Appendix C). In the next section we will combine the above result with the measure and write the complete expression for the two loop four-particle amplitude. Some properties of this chiral integrand will also be studied in the next section.

8 The 4-particle amplitude IV: the final result

The chiral integrand obtained in the last section is not symmetric under the interchange $q_1 \leftrightarrow q_2$. If we take the limit of $\tilde{p}_1 \rightarrow q_2$ we would obtain the

same result with $q_1 \leftrightarrow q_2$. So the final result should be a symmetrization of eq. (105). We suspect that this ambiguity is caused by our use of the delta function super Beltrami differentials and the final result actually requires this symmetrization if we study carefully the limit. We will not do this in this paper. In fact this symmetrization also makes the final amplitude explicitly independent of q as we will show immediately.

The symmetrized chiral integrand is

$$\begin{aligned} \mathcal{A} = & K(k_i, \epsilon_i) \langle : (\partial X(q_1) + \partial X(q_2)) \cdot (\partial X(q_1) + \partial X(q_2)) : \\ & \times \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \prod_{i=1}^4 \frac{q - z_i}{y(z_i)}. \end{aligned} \quad (106)$$

By using eq. (114) for the $\langle \partial X(z) \partial X(w) \rangle$ correlators we have

$$\langle : \partial X(q_1) \cdot (\partial X(q_1) + \partial X(q_2)) : \rangle = 0. \quad (107)$$

To compute explicitly eq. (106) we also need eq. (117) for the $\langle \partial X(z) X(w, \bar{w}) \rangle$ correlators and we have

$$\sum_{i=1}^4 \langle (\partial X^\mu(q_1) + \partial X^\mu(q_2)) k_i \cdot X(w_i) \rangle = - \sum_{i=1}^4 \frac{k_i^\mu}{q - w_i}. \quad (108)$$

By using eq. (107) and eq. (108) in eq. (106) we have:

$$\begin{aligned} \mathcal{A} = & K(k_i, \epsilon_i) \left[i \sum_{i=1}^4 \frac{k_i}{q - z_i} \right]^2 \prod_{i < j} \exp[-k_i \cdot k_j G(z_i, z_j)] \prod_{i=1}^4 \frac{q - z_i}{y(z_i)} \\ = & \frac{K(k_i, \epsilon_i)}{\prod_{i=1}^4 y(z_i)} \prod_{i < j} \exp[-k_i \cdot k_j G(z_i, z_j)] \\ & \times (s(z_1 z_2 + z_3 z_4) + t(z_1 z_4 + z_2 z_3) + u(z_1 z_3 + z_2 z_4)), \end{aligned} \quad (109)$$

which is independent of the insertion points $q_{1,2}$. Here $G(z_i, z_j)$ is the scalar Green function which is given in terms of the prime form $E(z_i, z_j)$ as follows (see [20]):

$$G(z, w) = -\ln |E(z, w)|^2 + 2\pi \text{Im} \int_z^w \omega_I (\text{Im} \Omega)_{IJ}^{-1} \text{Im} \int_z^w \omega_J. \quad (110)$$

s, t, u are the standard Mandelstam variables, $s = -(k_1 + k_2)^2$, etc.

For type II superstring theory the complete integrand is

$$\begin{aligned}
\mathcal{A}_{II} &= K(k_i, \epsilon_i) \langle : (\partial X(q_1) + \partial X(q_2)) \cdot (\partial X(q_1) + \partial X(q_2)) : \\
&\quad \times : (\bar{\partial} X(\tilde{q}_1) + \bar{\partial} X(\tilde{q}_2)) \cdot (\bar{\partial} X(\tilde{q}_1) + \bar{\partial} X(\tilde{q}_2)) : \\
&\quad \times \prod_{i=1}^4 e^{ik_i \cdot X(z_i, \bar{z}_i)} \rangle \prod_{i=1}^4 \frac{(q - z_i)(\tilde{q} - \bar{z}_i)}{|y(z_i)|^2} \\
&= \frac{K(k_i, \epsilon_i)}{\prod_{i=1}^4 |y(z_i)|^2} \prod_{i < j} \exp[-k_i \cdot k_j G(z_i, z_j)] \\
&\quad \times |s(z_1 z_2 + z_3 z_4) + t(z_1 z_4 + z_2 z_3) + u(z_1 z_3 + z_2 z_4)|^2, \quad (111)
\end{aligned}$$

which is independent of the left part insertion points $q_{1,2}$ and also the right part insertion points $\tilde{q}_{1,2}$.

The amplitude is obtained by integrating over the moduli space. At two loops, the moduli space can be parametrized either by the period matrix or three of the six branch points. We have

$$\begin{aligned}
A_{II} &= c_{II} K(k_i, \epsilon_i) \int \frac{\prod_{i=1}^6 d^2 a_i / dV_{pr}}{T^5 \prod_{i < j} |a_i - a_j|^2} \\
&\quad \times \prod_{i=1}^4 \frac{d^2 z_i}{|y(z_i)|^2} \prod_{i < j} \exp[-k_i \cdot k_j G(z_i, z_j)] \\
&\quad \times |s(z_1 z_2 + z_3 z_4) + t(z_1 z_4 + z_2 z_3) + u(z_1 z_3 + z_2 z_4)|^2, \quad (112)
\end{aligned}$$

where $dV_{pr} = \frac{d^2 a_i d^2 a_j d^2 a_k}{|a_{ij} a_{jk} a_{ki}|^2}$ is a projective invariant measure and c_{II} is a constant which should be determined by factorization or unitarity (of the S -matrix).

An immediate application of the above result is to study the perturbative correction to the R^4 term at two loops. By taking the limit of $k_i \rightarrow 0$, one sees from eq. (111) that the integrand $\mathcal{A}_{II}$ is zero identically in moduli space apart from the kinematic factor $K(k_i, \epsilon_i)$. So we conclude that the perturbative correction to the R^2 term is zero, confirming the explicit calculation of Iengo [10]. This result is also in agreement with the non-perturbative conjecture of [12] (see also [13, 14, 9]). Our new result also explicitly verifies the claim given in the the Appendix B of [15].

The finiteness of the amplitude can also be checked by following the detailed discussions given in [9].

Appendix A: The $\langle \partial X(z) \partial X(w) \rangle$ correlators

For the $X(z, \bar{z})$ correlators, D'Hoker and Phong in [21] use the following expression in terms of the prime form:

$$\langle \partial_z X^\mu(z) \partial_w X^\nu(w) \rangle = -\delta^{\mu\nu} \partial_z \partial_w \ln E(z, w), \quad (113)$$

in their effective rules for chiral splitting. In this paper we use the following result given in [29, 7]:

$$\begin{aligned} \langle \partial X^\mu(z) \partial X^\nu(w) \rangle = & -\delta^{\mu\nu} \left(\frac{1}{4(z-w)^2} + \frac{1}{4T} \frac{\partial}{\partial w} \left[\frac{y(w)}{y(z)} \frac{1}{z-w} \right. \right. \\ & \times \left. \left. \int \frac{(z-z_1)(z-z_2)}{(w-z_1)(w-z_2)} \left| \frac{z_1-z_2}{y(z_1)y(z_2)} \right|^2 d^2 z_1 d^2 z_2 \right] \right) + (z \leftrightarrow w). \end{aligned} \quad (114)$$

This correlator satisfies the following relation:

$$\int \bar{\Omega}_i(\bar{z}) \langle \partial X^\mu(z) \partial X^\nu(w) \rangle d^2 z = 0. \quad (115)$$

These two formulas are actually related by using eq. (110) by differentiating with respect to z and w . We have

$$\begin{aligned} \partial_z \partial_w \ln E(z, w) = & \pi \omega(z) \cdot (\text{Im} \tau)^{-1} \cdot \omega(w) \\ & + \left\{ \left(\frac{1}{4(z-w)^2} + \frac{1}{4T} \frac{\partial}{\partial w} \left[\frac{y(w)}{y(z)} \frac{1}{z-w} \right. \right. \right. \\ & \times \left. \left. \int \frac{(z-z_1)(z-z_2)}{(w-z_1)(w-z_2)} \left| \frac{z_1-z_2}{y(z_1)y(z_2)} \right|^2 d^2 z_1 d^2 z_2 \right] \right) + (z \leftrightarrow w) \right\}. \end{aligned} \quad (116)$$

By using eq. (114), one can also derive the correlators for $\sum_i \langle \partial X(z) k_i \cdot X(w_i) \rangle$ with $\sum_i \alpha_i = 0$. It is

$$\begin{aligned} \sum_i \langle \partial X^\mu(z) k_i \cdot X(w_i) \rangle = & -\sum_i k_i^\mu \left\{ \frac{1}{2(z-w_i)} + \frac{1}{2T} \int \frac{y(w_i)}{y(z)} \frac{1}{z-w_i} \right. \\ & \times \left. \frac{(z-z_1)(z-z_2)}{(w_i-z_1)(w_i-z_2)} \left| \frac{z_1-z_2}{y(z_1)y(z_2)} \right|^2 d^2 z_1 d^2 z_2 \right\}. \end{aligned} \quad (117)$$

The above correlator also satisfies

$$\int \bar{\Omega}_i(\bar{z}) \sum_i \langle \partial X^\mu(z) k_i \cdot X(w_i) \rangle d^2 z = 0. \quad (118)$$

By differentiating with respect to $\bar{w}_i$ in eq. (117), we have

$$\langle \partial X^\mu(z) \bar{\partial} X^\nu(\bar{w}) \rangle = -\delta^{\mu\nu} \left[\pi \delta^{(2)}(z - w) - \frac{\pi}{2} \omega(z) \cdot (\text{Im}\tau)^{-1} \cdot \bar{\omega}(\bar{w}) \right], \quad (119)$$

where

$$\begin{aligned} Q &\equiv \omega(z) \cdot (\text{Im}\tau)^{-1} \cdot \bar{\omega}(\bar{w}) \\ &= \frac{2}{T} \frac{1}{y(z)\bar{y}(\bar{w})} \int \frac{(z-u)(\bar{w}-\bar{u})}{|y(u)|^2} d^2u, \end{aligned} \quad (120)$$

and

$$T = \int d^2z_1 d^2z_2 \left| \frac{z_1 - z_2}{y(z_1)y(z_2)} \right|^2 = 2|\det K|^2 \det \text{Im}\tau. \quad (121)$$

The T_1 used in eq. (45) can also be obtained from eq. (114) explicitly. We have

$$\begin{aligned} T_1(w) &= \frac{1}{32} \left(\sum_{i=1}^6 \frac{1}{w - a_i} \right)^2 - \frac{1}{16} \sum_{i=1}^6 \frac{1}{(w - a_i)^2} \\ &\quad + \frac{1}{8T} \int d^2z_1 d^2z_2 \left| \frac{z_1 - z_2}{y(z_1)y(z_2)} \right|^2 \left[\frac{2}{(w - z_1)(w - z_2)} \right. \\ &\quad \left. + 2 \sum_{i=1}^2 \frac{1}{(w - z_i)^2} - \left(\frac{1}{w - z_1} + \frac{1}{w - z_2} \right) \sum_{i=1}^6 \frac{1}{w - a_i} \right]. \end{aligned} \quad (122)$$

The other formula used in the main text is

$$\partial_{p_a} \partial_{q_1} \ln E(p_a, q_1) + \partial_{p_a} \partial_{q_2} \ln E(p_a, q_2) = \frac{1}{(q - p_a)^2}, \quad (123)$$

which can be derived easily from eq. (116) by noting $y(q_2) = -y(q_1)$ and $\omega_i(q_2) = -\omega_i(q_1)$.

Appendix B: The proof of eq. (78)

The proof of the summation formulas is quite simple. The strategy is the same as the proof for other “vanishing identities” given in [28] which was first used in [6, 8, 7]. We first change $S_m(x)$ into a polynomial by using the following Möbius transformation:

$$a_i = x - \frac{1}{\tilde{a}_i}, \quad \text{or} \quad \tilde{a}_i = \frac{1}{x - a_i}, \quad \tilde{z}_i = \frac{1}{x - z_i}. \quad (124)$$

Noting that $y(z) = \prod_i^6 (z - a_i) = \prod_{i=1}^3 (z - A_i)(z - B_i)$, we can write $u(z)$ and $\frac{1}{u(z)}$ as follows:

$$u(z) = \frac{\prod_{i=1}^3 (z - A_i)}{y(z)}, \quad (125)$$

$$\frac{1}{u(z)} = \frac{\prod_{i=1}^3 (z - B_i)}{y(z)}. \quad (126)$$

By using these expressions we see that the left hand side of eq. (78) is a homogeneous polynomial in $\tilde{a}_i$ and $\tilde{z}_k$ (the Möbius transformed a_i and z_k) times $(\prod_{i=1}^4 \tilde{y}(\tilde{z}_k))^{-1}$, i.e.

$$\begin{aligned} P_{n,m} &= \sum_{\delta} \eta_{\delta} Q_{\delta} \left\{ \frac{u(z_1)u(z_2)}{u(z_3)u(z_4)} - (-1)^n \frac{u(z_1)u(z_2)}{u(z_3)u(z_4)} \right\} (S_m(x))^n \\ &= \frac{y^4(x)}{\prod_{i=1}^4 \tilde{y}(\tilde{z}_i)} \sum_{\delta} \eta_{\delta} \tilde{Q}_{\delta} \left[\prod_{i=1}^3 \prod_{j=1}^2 \prod_{l=3}^4 (\tilde{z}_k - \tilde{A}_i)(\tilde{z}_l - \tilde{B}_i) - (\tilde{A} \leftrightarrow \tilde{B}) \right] \\ &\quad \times \left(\sum_{i=1}^3 [(\tilde{A}_i)^m - (\tilde{B}_i)^m] \right)^n. \end{aligned} \quad (127)$$

This polynomial is modular invariant and it is also vanishing when $\tilde{z}_1 = \tilde{z}_{3,4}$ and $\tilde{z}_2 = \tilde{z}_{3,4}$. So it is proportional to $P(\tilde{a}) \prod_{i=1}^2 \prod_{j=3}^4 (\tilde{z}_i - \tilde{z}_j)$. The rest factor can only be obtained by explicit computation (most easily by computer). Then we have

$$P_{n,m} = \frac{y^4(x)}{\prod_{i=1}^4 \tilde{y}(\tilde{z}_i)} \times 2P(\tilde{a}) \prod_{i=1}^2 \prod_{j=3}^4 (\tilde{z}_i - \tilde{z}_j) \times C_{n,m}. \quad (128)$$

By an inverse Möbius transformation we have

$$\tilde{y}(\tilde{z}) = \prod_{i=1}^6 (\tilde{z} - \tilde{a}_i)^{1/2} = \frac{y(z)}{(x - z)^3 y(x)}, \quad (129)$$

$$P(\tilde{a}) = \frac{P(a)}{(y(x))^{10}}, \quad (130)$$

etc. By using these results in eq. (128), we obtained the results given in eqs. (79)–(82). This completes the proof of eq. (78).

Appendix C: The q dependence of the factor $\mathcal{Z}$

In this appendix we will compute explicitly the q dependence of the factor $\mathcal{Z}$ defined in eq. (34). The relevant part is

$$\int \mathcal{D}\beta \mathcal{D}\gamma \delta(\beta(q_1)) \delta(\beta(q_2)) e^{iS_{\beta\gamma}} = (\det' \bar{\partial}_{3/2})^{-1} \int d\beta_1 d\beta_2 \delta(\beta_0(q_1)) \delta_0(\beta(q_2)), \quad (131)$$

where β_1 and β_2 are the two coefficients in the zero mode part of the ghost field $\beta(z)$:

$$\beta_0(z) = \beta_1 \varphi_1^*(z) + \beta_2 \varphi_2^*(z), \quad (132)$$

where $\varphi_{1,2}^*(z)$ are two holomorphic $\frac{3}{2}$ -differentials. Comparing with the previously defined two holomorphic $\frac{3}{2}$ -differential $\psi_{1,2}^*(z)$, $\varphi_{1,2}^*(z)$ are not required to satisfy the “normalization” condition eq. (22).

Choosing an arbitrary 1-differential $\omega_x(z)$ which has two zeroes at $z = x_1 = x_+$ and $z = x_2 = x_-$, the $\varphi_{1,2}^*(z)$ can be constructed as follows:

$$\varphi_\alpha^*(z) = \omega_x(z) S(z, x_\alpha). \quad (133)$$

By using these holomorphic differentials, we have

$$\begin{aligned} & \int d\beta_1 d\beta_2 \delta(\beta_0(q_1)) \delta(\beta_0(q_2)) \\ &= \int d\beta_1 d\beta_2 \prod_{\alpha=1}^2 \delta(\beta_1 \omega_x(q_\alpha) S(q_\alpha, x_1) + \beta_2 \omega_x(q_\alpha) S(q_\alpha, x_2)) \\ &= |\det(\omega_x(q_\alpha) S(q_\alpha, x_\beta))|^{-1} = \left[\frac{q-x}{\omega_x(q)} \right]^2, \end{aligned} \quad (134)$$

by using the explicit expression of the Szegő kernel and noting that $\omega_x(q_2) = -\omega_x(q_1)$ and $u(q_2) = -u(q_1)$, etc. In hyperelliptic language, the 1-differential $\omega_x(z)$ is given as follows:

$$\omega_x(z) = \frac{c(z-x)}{y(z)}, \quad (135)$$

where c is a possibly moduli dependent constant. By using this result we have:

$$\int \mathcal{D}\beta \mathcal{D}\gamma \delta(\beta(q_1)) \delta(\beta(q_2)) e^{iS_{\beta\gamma}} = \langle \delta(\beta(q_1)) \delta(\beta(q_2)) \rangle = \frac{y^2(q)}{c^2} \times (\det' \bar{\partial}_{3/2})^{-1}. \quad (136)$$

The q dependent factor $y^2(q)$ exactly cancels the factor $y^2(q)$ in $\mathcal{A}$. The complete expression for $\mathcal{Z}$ is:

$$\mathcal{Z} = \frac{y^2(q) \prod_{i < j} A_{ij} B_{ij}}{c^2 (\det K)^5 \prod_{i < j} (a_i - a_j)} = \frac{y^2(q) Q_\delta}{c^2 (\det K)^5 P(a)}, \quad (137)$$

by using the results of [8] (for previous works see [40, 41, 42, 43, 44, 45]).

Finally we have a variational formula from [32]:

$$\frac{\partial \tau_{ij}}{\partial a_n} = \frac{i\pi}{2} \hat{\omega}_i(a_n) \hat{\omega}_j(a_n), \quad (138)$$

where $\omega(a_n)$ is defined as follows:

$$\begin{aligned} \omega(z) dz &= (\hat{\omega}(z_0) + \hat{\omega}'(z_0)(z - z_0) + \dots) dz \\ &= 2u\omega(u^2 + a_n) du, \end{aligned} \quad (139)$$

$$\omega(a_n) = \lim_{u \rightarrow 0} 2u\omega(u^2 + a_n). \quad (140)$$

Here we have used the uniformization coordinate u instead of z at the branch point a_n .

By using eq. (138), we have:

$$\frac{\partial(\tau_{ij})}{\partial(a_1 a_2 a_3)} = i \left(\frac{\pi}{2} \right)^3 \frac{a_{45} a_{46} a_{56}}{(\det K)^3 P(a)}, \quad (141)$$

which can be used to transform from period matrix parametrization to the branch points parametrization of the moduli space.

Acknowledgments

Chuan-Jie Zhu would like to thank Roberto Iengo for reading the paper and comments. He would also like to thank E. D'Hoker and D. Phong for discussions and Jian-Xin Lu and the hospitality at the Interdisciplinary Center for Theoretical Study, Physics, University of Science and Technology of China.

References

- [1] M. B. Green and J. H. Schwarz, ‘‘Supersymmetric Dual String Theory III’’, Nucl. Phys. **B198** (1982) 441.

- [2] J. H. Schwarz, “Superstring Theory”, Phys. Reports **89** (1982) 223–322.
- [3] M. B. Green, “Lectures on Superstrings”, in *Unified String Theories*, edited by M. Green and D. Gross (World Scientific, 1986).
- [4] E. Martinec, “Nonrenormalization Theorems and Fermionic String Finiteness”, Phys. Lett. **171B** (1986) 189–194.
- [5] E. Gava, R. Iengo and G. Sotkov, “Modular Invariance and the Two-Loop Vanishing of the Cosmological Constant”, Phys. Lett. **207B** (1988) 283–291.
- [6] R. Iengo and C.-J. Zhu, “Notes on the Non-Renormalization Theorem in Superstring Theories”, Phys. Lett. **212B** (1988) 309–312.
- [7] C.-J. Zhu, “Two-Loop Computations in Superstring Theories”, Int. J. Mod. Phys. **A4** (1989) 3877–3906.
- [8] R. Iengo and C.-J. Zhu, “Two-Loop Computation of the Four-Particle Amplitude in Heterotic String Theory”, Phys. Lett. **212B** (1988) 313–319.
- [9] R. Iengo and C.-J. Zhu, “Explicit Modular Invariant Two-Loop Superstring Amplitude Relevant for R^4 ”, J. High Energy Phys. **0006** (1999) 011, hep-th/9905050.
- [10] R. Iengo, “Computing the R^4 Term at Two Superstring Loops”, J. High Energy Phys. **0202** (2002) 035, hep-th/0202058.
- [11] D. J. Gross and E. Witten, “Superstring Modification of Einstein’s Equations”, Nucl. Phys. **B277** (1986) 1.
- [12] M. B. Green and M. Gutperle, “Effects of D-instantons”, Nucl. Phys. **B498** (1997) 195–227, hep-th/9701093.
- [13] M. B. Green, M. Gutperle and P. Vanhove, “One Loop in Eleven Dimension”, Phys. Lett. **409B** (1997) 277–184.
- [14] M. B. Green and S. Sethi, “Supersymmetry Constraints on Type IIB Supergravity”, Phys.Rev. **D59** (1999) 046006, hep-th/9808061.

- [15] S. Stieberger and T.R. Taylor, “Non-Abelian Born-Infeld action and type I - heterotic duality (I): Heterotic F^6 terms at two loops”, Nucl.Phys. **B647** (2002) 49–68, hep-th/0207026; “Non-Abelian Born-Infeld Action and Type I - Heterotic Duality (II): Nonrenormalization Theorems”, Nucl. Phys. **B648** (2003) 3, hep-th/0209064.
- [16] M. B. Green, “Interconnections Between Type II Superstrings, M Theory and N=4 Supersymmetric Yang–Mills” , hep-th/9903124
- [17] E. Kiritsis, “Duality and Instantons in String Theory”, in *Proceedings of the 1999 Spring Workshop on Superstrings and Related Matters*, eds. B. Greene, J. Louis, K.S. Narain and S. Randjbar-Daemi (World Scientific, Singapore, 2000), hep-th/9906018
- [18] E. D’Hoker and D.H. Phong, “The Geometry of String Perturbation Theory”, Rev. Mod. Phys. **60** (1988) 917–1065.
- [19] E. D’Hoker and D.H. Phong, “Lectures on Two-Loop Superstrings—Hangzhou, Beijing 2002”, hep-th/0211111.
- [20] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings I, Main Formulas”, Phys. Lett. **B529** (2002) 241–255.
- [21] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings II, The chiral Measure on Moduli Space”, Nucl. Phys. **B636** (2002) 3–60.
- [22] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings III, Slice Independence and Absence of Ambiguities”, Nucl. Phys. **B636** (2002) 61–79.
- [23] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings IV, The Cosmological Constant and Modular Forms”, Nucl. Phys. **B639** (2002) 129–181.
- [24] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings V, Scattering Amplitudes, the 1-, 2- and 3-point functions”, in preparation.
- [25] E. D’Hoker and D.H. Phong, “Two-Loop Superstrings VI, Scattering Amplitudes: the 4-point function”, in preparation.
- [26] O. Lechtenfeld and A. Parkes, “On Covariant Multiloop Superstring Amplitudes”, Nucl. Phys. **B332** (1990) 39–82.

- [27] Zhu-Jun Zheng, Jun-Bao Wu and C.-J. Zhu, “Two-Loop Superstrings in Hyperelliptic Language I: the Main Results”, hep-th/0212191.
- [28] Zhu-Jun Zheng, Jun-Bao Wu and C.-J. Zhu, “Two-Loop Superstrings in Hyperelliptic Language II: the Vanishing of the Cosmological Constant and the Non-Renormalization Theorem”, hep-th/0212198.
- [29] V. G. Knizhnik, “Explicit Expression for the Two-Loop Measure in the Heterotic String Theory”, Phys. Lett. **196B** (1987) 473–476.
- [30] A. Morozov, “Two-Loop Statsum of Superstrings”, Nucl. Phys. **B303** (1988) 343–372; “On the Two-Loop Contribution to the Superstring Four-Point Function”, Phys. Lett. **209B** (1988) 473–476.
- [31] A. Morozov and A. Pereomov, “Partition Functions in Superstring Theory. The Case of Genus Two”, Phys. Lett. **197B** (1987) 115–118; “A Note on Many-Loop Calculations for Superstrings in the NSR Formalism”, Int. J. Mod. Phys. **A4** (1989) 1773–1780.
- [32] D. Lebedev and A. Morozov, “Statistical Sums of Strings on Hyperelliptic Riemann Surfaces”, Nucl. Phys. **B302** (1988) 163–188.
- [33] O. Lechtenfeld and A. Parkes, “On the Vanishing of the Genus Two Superstring Vacuum Amplitude”, Phys. Lett. **202B** (1988) 75–80.
- [34] M. A. Bershadsky, “The Superbranch Point and Two-Loop Correlation in Heterotic String Theory”, Mod. Phys. Lett. **A3** (1988) 91–108.
- [35] G. Moore and A. Morozov, “Some Remarks on Two-Loop Superstring Calculations”, Nucl. Phys. **B306** (1988) 387–404.
- [36] E. Verlinde and H. Verlinde, “Multiloop Calculations in Covariant Superstring Theory”, Phys. Lett. **192B** (1987) 95–102.
- [37] M. Bonini and R. Iengo, “Correlation Functions and Zero Modes on Higher Genus Riemann Surface”, Int. J. Mod. Phys. **A3** (1988) 841–860.
- [38] J. J. Atick and A. Sen, “Covariant One-Loop Fermion Amplitudes in Closed String Theories”, Nucl. Phys. **B293** (1987) 317–347.

- [39] C.-J. Zhu, “Covariant Two-Loop Fermion Amplitude in Closed Superstring Theories”, Nucl. Phys. **B327** (1989) 744–762.
- [40] L. Dixon, D. Friedan, E. Martinec and S. Shenker, “The Conformal Field Theory of Orbifolds”, Nucl. Phys. **B282** (1987) 13–73.
- [41] Al. B. Zamolodchikov, “Conform Scalar Field on the Hyperelliptic Curves and Critical Ashkin-Teller Multipoints Correlation Functions”, Nucl. Phys. **B285** [**FS19**] (1987) 481–503.
- [42] M. A. Bershadsky and A. O. Radul, “Conformal Field Theories with Additional $\mathbf{Z}_N$ Symmetry”, Int. J. Mod. Phys. **A2** (1987) 165–178; “ g -Loop Amplitudes in Bosonic String Theory in Terms of Branch Points”, Phys. Lett. **193B** (1987) 213–218.
- [43] V. G. Knizhnik, “Analytic Fields on Riemann Surface. II”, Comm. Math. Phys. **112** (1987) 567–590.
- [44] D. Montano, “Superstrings on Hyperelliptic Surfaces and the Two-Loop Vanishing of the Cosmological Constant”, Nucl. Phys. **B297** (1988) 125–140.
- [45] E. Gava, “Conformal Fields on Complex Curves”, preprint LEPTENS 88/1 (Jan., 1988) 15 pages.